

General Mathematics Review, Linear Algebra Overview, and Using Python for Linear Algebra

Julie Butler

Disclaimer: You will not have to do *much* math by hand in this course, but it is good to know how to do some basic operations and what the terminology means.

Scalars

Scalars

- ▶ A scalar is just a number, what we are use to dealing with in math.
- ▶ Scalars can have a decimal place or be a whole number (called an integer).
- ▶ Scalars can be greater than zero (positive) or less than zero (negative)
- ▶ Examples:
 - ▶ 5
 - ▶ 6.7292
 - ▶ -8.1

Arithmetic with Scalars

- ▶ Scalars can have four basic arithmetic operators performed on them: addition, subtraction, multiplication, and division.
 - ▶ Order does not matter for addition and multiplication, but does matter for subtraction and division.
- ▶ Example: Give $a = 5$ and $b = 2$, find:
 - ▶ $a+b$
 - ▶ $a-b$
 - ▶ $b-a$
 - ▶ ab ($a * b$)
 - ▶ a/b
 - ▶ b/a

Factoring and Distribution

- ▶ Consider the scalars a , b , c :
 - ▶ If they are arranged as $ab+ac$, then since both terms of the sum have the same scalar (a) you can *factor* it out: $a(b+c)$
 - ▶ Implicit multiplication
 - ▶ If they are arranged as $(a+b)(a-c)$ then we can *distribute* the first part of the term $(a+b)$ to both parts of the second term.
 - ▶ $(a+b)a - (a+b)c$
 - ▶ We can then *distribute* the a or the c to both scalars in parentheses
 - ▶ $aa + ab - ac + bc = a^2 + ab - ac + bc$ (exponents)
 - ▶ Note that we could also have used FOIL here (First, Outer, Inner, Last)

Scalars in Python

```
# Defining scalars (as floats, not ints)
a = 5.0
b = 4.0

# Addition, Subtraction, Multiplication, Division
add = a+b
sub = a-b
mul = a*b
div = a/b
print("Scalar Math:", add, sub, mul, div)

# Multiplication with Parentheses
# Multiplication is not implicit.
mul_paren = (a+b)*(a-b)
print("Multiplication with Partentheses:", mul_paren)
```

Complex Numbers

Complex Numbers

- ▶ A complex number has both a real part (a) and an imaginary part (b): $z = a+ib$
 - ▶ i : the imaginary number $i = \sqrt{-1}$
 - ▶ Note that the imaginary part is just b , not ib .
- ▶ Example: Find the real and imaginary parts of the following numbers.
 - ▶ $z = 2+3i$
 - ▶ $z = -5i$
 - ▶ $z = 7$

Arithmetic with Complex Numbers

- ▶ Consider the complex numbers $x = a+ib$ and $y = c+id$
 - ▶ Addition and Subtraction: $x \pm y = (a \pm c) + i(b \pm d)$
 - ▶ Add/subtract the real parts and then add/subtract the imaginary parts
 - ▶ Multiplication: $xy = (a+ib)(c+id) = ac + iad + ibc + i(i)bd = (ac-bd) + i(ad+bc)$
 - ▶ FOIL two imaginary numbers to multiply
 - ▶ Note that $i * i = \sqrt{-1}\sqrt{-1} = (-1)^2 = -1$
 - ▶ Example: If $x = 2-3i$ and $y = 4+i$, find $x+y$, $x-y$, and xy .

Complex Conjugate

- ▶ The complex conjugate of a complex number z is denoted as z^* and is found by negating the imaginary part of z
- ▶ If $z = a+ib$ then $z^* = a-ib$
- ▶ Example: If $z = 5+2i$, find z^* .

Norm or Modulus

- ▶ The modulus squared of a complex number (denoted as $|z|^2$) is just the number times its complex conjugate.
- ▶ If $z = a+ib$ then
$$zz^* = |z|^2 = (a + ib)(a - ib) = a^2 - iab + iab - i(i)b^2 = a^2 + b^2$$
- ▶ The modulus (or norm) is just the square root of the modulus squared: $|z| = \sqrt{a^2 + b^2}$
- ▶ The modulus and modulus squared of a complex number is always a real number.
- ▶ Example: If $z = 4+3i$, find the modulus and the modulus squared.

Complex Numbers in Python

- ▶ Complex numbers are built into Python, but you denote the imaginary number, i , with a j instead
- ▶ Note that because we defined the numbers to be complex, it does not remove the imaginary term, even when it's zero

```
import numpy as np

# Define a complex number
z = 4+3j
print("Complex Number:", z)

# Find the complex conjugate
z_conj = z.conjugate()
print("Complex Conjugate:", z_conj)

# Find the modulus and modulus squared
# Note that the modulus and modulus squared are still
# complex numbers
z_mod_squared = z*z_conj
z_mod = np.sqrt(z_mod_squared)
print("Modulus Squared and Modulus:", z_mod_squared, z_mod)

# Basic Arithmetic
x = 1+2j
y = 3-4j
print("Complex Number Math:", x+y, x-y, x*y)
```

Vectors, Bras, Kets (Dirac Notation)

Vectors

- ▶ A vector is a one-dimensional structure of numbers arranged in either a row or a column.

$$\vec{a} = [1 \quad 2 \quad 3]$$

$$\vec{b} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

- ▶ Each number in a vector is called an *element*, the number of elements is the size/length/rank of the vector
 - ▶ Elements can be real or imaginary

Magnitude and Unit Vectors

- ▶ A vector has a size and a direction.
- ▶ The size of a vector is called its magnitude, calculated by summing the square of all the elements then taking the square root.
 - ▶ Example: $\vec{a} = [1 \ 2 \ 3]$. Find a (also denoted as $|\vec{a}|$), the magnitude of \vec{a}
- ▶ The direction of a vector is denoted by its unit vector, found by dividing the vector by its magnitude.
 - ▶ Example: $\vec{a} = [1 \ 2 \ 3]$. Find \hat{a} , the unit vector of \vec{a} .
 - ▶ Note that the magnitude of a unit vector is 1.
- ▶ Note that $|\vec{a}|\hat{a} = \vec{a}$

Arithmetic with Vectors

- ▶ If you want to add or subtract two vectors, just add or subtract their corresponding components:
- ▶ Example: $\vec{a} = [1 \ 2 \ 3]$ and $\vec{b} = [4 \ 5 \ 6]$. Find $a+b$ and $a-b$.
- ▶ The vectors must be the same length to be added or subtracted.
- ▶ You can multiply or divide a vector by a scalar if you perform that operation on every element of the vector.
- ▶ Example: $\vec{a} = [1 \ 2 \ 3]$ and $c = 10$. Find ca and a/c .
- ▶ Note that you **cannot** divide by a vector.

Dot Products

- ▶ There are two ways to “multiply” vectors; the first is called the dot product (also called the inner product and the scalar product)
- ▶ Multiple elements in like locations and then add them all together
 - ▶ Example: $\vec{a} = [1 \ 2 \ 3]$ and $\vec{b} = [4 \ 5 \ 6]$. Find $\vec{a} \cdot \vec{b}$.
 - ▶ Note that the result of calculating a dot product is a **scalar**
- ▶ Note that $\vec{a} \cdot \vec{b} = |\vec{a}||\vec{b}|\cos\theta$, where θ is the angle between the two vectors

Cross Products

- ▶ The other way to “multiply” vectors is using a cross product, these are a bit complicated to calculate by hand so we won't do that in this course.
 - ▶ However, note that $|\vec{a} \times \vec{b}| = |\vec{a}||\vec{b}|\sin\theta$.
- ▶ A cross product will produce a **vector** which is perpendicular to both vectors.

Bras and Kets (Dirac Notation)

- ▶ In mathematics, vectors will be represented as a, \mathbf{a} , or \vec{a} , almost always with a lowercase letter.
- ▶ In quantum mechanics, we use a slightly different notation
 - ▶ Greek letters: $\alpha, \beta, \phi, \psi$
 - ▶ Ket: $|\psi\rangle \rightarrow$ always a column vector
 - ▶ Bra: $\langle\psi| \rightarrow$ always a row vector
- ▶ A ket can be transitioned into a bra by turning it into a row vector and taking the complex conjugate of each element:
 - ▶ Example:

$$|\psi\rangle = \begin{bmatrix} 2 - 5i \\ 6i \\ 3 \end{bmatrix}$$

$$\langle\psi| = [2 + 5i \quad -6i \quad 3]$$

Dirac Notation Continued

- ▶ A bra and a ket written together form a bracket: $\langle \phi | \psi \rangle$
 - ▶ A bracket is equivalent to performing an inner product (dot product)
- ▶ Example:

$$\langle \phi | = [1 \quad 2 \quad 3]$$

$$|\psi\rangle = \begin{bmatrix} 2 - 5i \\ 6i \\ 3 \end{bmatrix}$$

$$\langle \phi | \psi \rangle = 1(2 - 5i) + 2(6i) + 3(3) = 2 - 5i + 12i + 9 = 7i + 11$$

- ▶ Note that we typically what bras and kets to have a magnitude of 1, so $\langle \psi | \psi \rangle = 1$

Vectors in Python

```
import numpy as np

# Define vectors as Numpy arrays
a = np.array([1,2,3])
b = np.array([4,5,6])
c = 10

# Arithmetic with Vectors
add = a+b
sub = a-b
mul = c*a
print("Vector Math:", add, sub, mul)

# Dot product two ways
dot1 = np.dot(a,b) #Order is important
dot2 = a.dot(b) # Order is important
print("Dot Product:", dot1, dot2)

# Cross product
cross = np.cross(a,b)
print("Cross Product:", cross)

# Magnitude and unit vector
mag = np.linalg.norm(a)
unit_vector = a/mag
unit vector mag = np.linalg.norm(unit vector)
```



Matrices

Matrices

- ▶ Matrices are two-dimensional structures where each *element* can be a real or imaginary number
 - ▶ The width and height can be the same (square matrix) or different. We will mostly deal with square matrices
- ▶ Example:

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

Arithmetic with Matrices

- ▶ Matrices add and subtract just like vectors (do the operations to like elements), and can be multiplied or divided by a scalar like vectors.
- ▶ Example: Given the following matrices and $c = 10$, find $A+B$, $A-B$, cA and A/c .

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

$$B = \begin{bmatrix} 9 & 8 & 7 \\ 6 & 5 & 4 \\ 3 & 2 & 1 \end{bmatrix}$$

Matrix Multiplication and Commutator

- ▶ Two matrices can be multiplied together but this process is tedious to do by hand so we will use Python or other solvers
- ▶ In general, $AB \neq BA$, or phrased another way, $AB - BA \neq 0$
- ▶ We can define the commutator of two matrices: $[A,B] = AB - BA$
 - ▶ if $[A,B] = 0$ (meaning $AB = BA$) we say that A and B commute

Arithmetic with Matrices and Vectors

- ▶ You cannot add or subtract vectors or matrices, but you can multiply a vector and a matrix. The result will be a vector
- ▶ This again is a bit tedious to do by hand so we will use Python or other solvers

Transpose, Adjoint, Conjugate, and More

- ▶ Consider the matrix

$$A = \begin{bmatrix} 1 & 2i & 3 \\ 4+i & 5 & 6 \\ 7 & 8 & 9-3i \end{bmatrix}$$

- ▶ The transpose of A , A^T , is found by switching the rows and columns.
- ▶ The complex conjugate of A , A^* is found by taking the complex conjugate of every element.
- ▶ The adjoint of a matrix, $(A^T)^* = A^\dagger$ is found by taking the transpose **and** the complex conjugate of a matrix. If $A = A^\dagger$ then the matrix is said to be *Hermitian*

Dirac Notation

- ▶ Matrices are not represented differently in Dirac notation.
- ▶ $A|\psi\rangle$ is a matrix multiplied by a vector (gives a vector)
- ▶ $\langle\phi|A|\psi\rangle$ is a vector multiplied by a matrix multiplied by a vector

Eigenvalues and Eigenvectors

- ▶ Eigenvalues (scalars) and eigenvectors are special values and vectors that each matrix has
 - ▶ The eigenvalue, λ and the eigenvector, x , solve the equation
$$Ax = \lambda x$$
- ▶ You can think of eigenvalues as the “roots” of the matrix
- ▶ Finding these by hand can be complicated, we will use Python or other solvers

Matrices in Python

```
import numpy as np
# Matrices are defined as two-dimensional Numpy arrays
A = np.array([[1,2,3], [4,5,6], [7,8,9]])
B = np.array([[9,8,7], [6,5,4], [3,2,1]])
# Define a vector
x = np.array([2,4,6])
# Define a scalar
c = 10

# Matrix addition and subtraction
add = A+B
sub = A-B
print("Matrix Addition:", add)
print("Matrix Subtraction:", sub)
print()

# Matrix-matrix multiplication
# Do not use the * symbol!!
mul1 = A@B
mul2 = B@A
commutator = mul1 - mul2
print("Matrix Multiplication 1:",mul1)
print("Matrix Multiplication 2:", mul2)
print("Commutator:", commutator)
print()

# Matrix-Vector Multiplication
mul3 = A@x
print("Matrix-Vector Multiplication:", mul3)
print()

# Matrix-Scalar Multiplication
# Use the * symbol!!
mul4 = c*B
print("Matrix-Scalar Multiplication:", mul4)
print()

# Transpose and Adjoint
transpose = A.T
adjoint = np.conj(B).T
print("Transpose:", transpose)
print("Adjoint:", adjoint)
print()

# Eigenvalues and Eigenvectors
# Note that the eigenvectors of A are the columns of the matrix
# eigenvectors
eigenvalues, eigenvectors = np.linalg.eig(A)
print("Eigenvalues:",eigenvalues)
print("Eigenvectors:",eigenvectors)
```